



Account #	Date:

COOLSCULPTING

POST CARE TREATMENT INSTRUCTIONS & COMMON SIDE EFFECTS

Post Care

- Tylenol can be taken for any discomfort during the healing process
- Avoid NSAIDs: such as Ibuprofen, Aspirin, Motrin, Advil, Aleve for 2 weeks post treatment. These over the counter medications interrupt the desired effects of the treatment.
- Over-the-counter antihistamines, like Benadryl, can be used for itching
- ❖ Hot/cold packs can be applied to the area to help with any discomfort
- Compression garments may be used on the treated area to add additional support and provide comfort during the healing process
- ❖ Avoid Turmeric for 2 weeks post treatment

Possible Side Effects

- Numbness and/or tingling (pins & needles sensation) may be present for up to 2-3 weeks in the treatment area after treatment and will resolve on its own.
- Swelling for up to 2-3 weeks in the treatment area is common and will resolve
- Bruising in the treatment area may occur due to the suction of the applicators applied during treatment
- Itching in the treatment area can also be present for up to 2-3 weeks post treatment
- Side effects may vary for individuals. Please contact our office if your side effects persist more than 3 weeks.
- Visible results can take up to 4 months

A 90 day follow-up appointment will be scheduled after your initial treatment to take additional photos and evaluate the results of the treatment up to that point.

<u>Please call our office if you have any questions or concerns regarding your treatment at</u> 480-556-0446.

Scottsdale . Surprise . Anthem . Deer Valley . Gilbert . Ahwatukee . Tempe